



In childhood diarrheas

- careful supervision
- electrolyte replacement
- specific anti-infective therapy and

LOMOTIL®

tablets/liquid

Each tablet and each 5 cc. of liquid contains:
 diphenoxylate hydrochloride 2.5 mg.
 (Warning: May be habit forming)
 atropine sulfate 0.025 mg.

Warnings: Lomotil should be used with caution in patients taking barbiturates and with caution, if not contraindicated, in patients with cirrhosis, advanced liver disease or impaired liver function.

Precautions: Lomotil is a Federally exempt narcotic with theoretically possible addictive potential at high dosage; this is not ordinarily a clinical problem. Use Lomotil with considerable caution in patients receiving addicting drugs. Recommended dosages should not be exceeded, and medication should be kept out of reach of children. Should accidental overdose occur, signs may include severe respiratory depression, flushing, lethargy or coma, hypotonic reflexes, nystagmus, pinpoint pupils, tachycardia; continuous observation is necessary.

Adverse Reactions: Side effects reported with Lomotil therapy include nausea, sedation, dizziness, vomiting, pruritus, restlessness, abdominal discomfort, headache, angioneurotic edema, giant urticaria, lethargy, anorexia, numbness of the extremities, atropine effects, swelling of the gums, euphoria, depression and malaise. Respiratory depression and coma may occur with overdose.

LOMOTIL in conjunction with specifically indicated medical management may be life saving in children with severe diarrhea associated with such conditions as acute infections, gastroenteritis, drug therapy and food poisoning.

Lomotil lowers the excessive intestinal propulsion characteristic of diarrhea. This reduction of precipitate intestinal flow allows a normal or more nearly normal reabsorption of fluid and electrolytes and counteracts the dehydration so hazardous to children.

This specific, well localized pharmacologic activity controls both acute infectious diarrheas and long-term functional and organic diarrhea with unsurpassed promptness, convenience and efficiency.

PROMPT • EFFECTIVE • CONVENIENT

Dosage: The recommended initial daily dosages, given in divided doses until diarrhea is controlled, are as follows:

Children: Total Daily Dosage

3-6 mo. . . . ½ tsp.* t.i.d. (3 mg.) ↓ ↓ ↓
 6-12 mo. . . ½ tsp. q.i.d. (4 mg.) ↓ ↓ ↓ ↓
 1-2 yr. . . . ½ tsp. 5 times daily (5 mg.) ↓ ↓ ↓ ↓ ↓
 2-5 yr. . . . 1 tsp. t.i.d. (6 mg.) ↓ ↓ ↓
 5-8 yr. . . . 1 tsp. q.i.d. (8 mg.) ↓ ↓ ↓ ↓
 8-12 yr. . . 1 tsp. 5 times daily (10 mg.) ↓ ↓ ↓ ↓ ↓
Adults: . . . 2 tsp. 5 times daily (20 mg.) ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓
 or 2 tablets q.i.d. ○ ○ ○ ○ ○ ○ ○ ○

*Based on 4 cc. per teaspoonful.

Maintenance dosage may be as low as one-fourth the initial daily dosage.

SEARLE

*Research in the
Service of Medicine*

You're the reason we're here.

You and your patients are the reason we got started, the reason we grew up, and the reason we want to keep moving ahead.

That's also the reason we have practicing physicians on our committees. Like the Physicians' Review Committee, which acts as your own personal court of appeals when you have a question about a Blue Shield allowance. We're glad this committee is made up of

doctors because if anyone is aware of your problems, they are.

And we want you to be aware of our policies and to let us know how they affect you. That's why you have direct lines to us through the Medical Societies' Reference Committee and your Professional Relations representatives. The Committee works through your local Medical Society, and the representatives contact you directly. They both help to make Blue Shield more aware of your needs even though you don't have time to sit in on meetings

and planning committees.

We'd like to feel that when we represent you and your interests on both the local and the national levels, we're representing our own. But if you know what we can do to bring our interests and yours closer together, let us know. If it will make Blue Shield better for you and your patients, that's a good enough reason for us.



® GREATER NEW YORK'S
BLUE SHIELD
UNITED MEDICAL SERVICE, INC.

3

Polymyxin B—Bacitracin—Neomycin

against 10



Pseudomonas




Hemophilus



Klebsiella



Aerobacter




Escherichia



Proteus




Corynebacterium



Staphylococcus



Streptococcus



Pneumococcus

‘Neosporin’[®] Ointment Polymyxin B—Bacitracin—Neomycin

Overlapping, broad bactericidal coverage.

Nonirritant ointment base; also enhances spreading and penetration.

Each gram contains:

‘Aerosporin’[®]

brand Polymyxin B Sulfate 5,000 Units

Zinc Bacitracin 400 Units

Neomycin Sulfate 5 mg.
(equivalent to 3.5 mg. Neomycin Base)

Special White Petrolatum q.s.

Contraindications: This product is contraindicated in those individuals who have shown hypersensitivity to any of its components. Do not use in the external ear canal if the eardrum is perforated.

Precautions: As with other antibiotic products,


prolonged use may result in overgrowth of non-susceptible organisms, including fungi. Appropriate measures should be taken if this occurs. Articles in the current medical literature indicate an increase in the prevalence of persons allergic to neomycin. The possibility of such a reaction should be borne in mind.

Available: Tubes of 1 oz., 1/2 oz. with applicator tip, 1/8 oz. with ophthalmic tip. The ointment base and the formula of the various sizes are identical, but only the 1/8 oz. tube should be used for ophthalmic purposes.



BURROUGHS WELLCOME & CO. (U.S.A.) INC., Tuckahoe, New York

now for your non-hospitalized patients



Serum Chemistry Graph provided is 8" x 11"

blood chemistry profiles

run on
TECHNICON®
Sequential Multiple Analyzer

only \$15⁰⁰

FOR A ~~12~~-TEST BLOOD CHEMISTRY PROFILE

Heretofore available only to patients in a few large hospitals, PATHOLOGICAL LABORATORY now provides this rapid automated analytical service for the non-hospitalized patient.

A Serum Chemistry Graph of each analysis summarizes in graphic form the results of all 12 tests together, superimposed over a background of normal values.

Patients may be directed to the laboratory; or blood samples can be mailed in post-paid containers provided by our laboratory. The Serum Chemistry Graph and a report in the usual numerical format will be mailed the same day samples are received. Free messenger pick-up is available in select areas.

PATHOLOGICAL LABORATORY

40-03 NATIONAL ST., CORONA, N.Y. 11368

TELEPHONE 212-639-5403

**send for
post-paid
blood sample
mailing
containers**

PATHOLOGICAL LABORATORY • 40-03 National St., Corona, N.Y. 11368

Gentlemen:


- ☐ Please send additional literature
- ☐ Please send post-paid containers for blood samples

Dr.

Street

City State Zip

It's almost as if you were there to give an injection of penicillin



V-Cillin K[®], Pediatric dependable oral penicillin therapy

Potassium Phenoxymethyl Penicillin

Description: V-Cillin K, the potassium salt of V-Cillin[®] (phenoxymethyl penicillin, Lilly), combines acid stability with immediate solubility and rapid absorption. Higher, more rapid serum levels are obtained than with equal oral doses of penicillin G.

Indications: Streptococcus, pneumococcus, and gonococcus infections; infections caused by sensitive strains of staphylococci; prophylaxis of streptococcus infections in patients with a history of rheumatic fever; and prevention of bacterial endocarditis after tonsillectomy and tooth extraction in patients with a history of rheumatic fever or congenital heart disease.

Contraindication: Penicillin hypersensitivity.

Warnings: In rare instances, penicillin may cause acute anaphylaxis which may prove fatal unless promptly controlled. This type of reaction appears more frequently in patients with a history of sensitivity reactions to penicillin or with bronchial asthma or other allergies. Resuscitative drugs should be readily available. These include epinephrine and pressor drugs (as well as oxygen for inhalation) for immediate allergic manifestations and antihistamines and corticosteroids for delayed effects.

Precautions: Use cautiously, if at all, in a patient with a strongly positive history of allergy.

In prolonged therapy with penicillin, and particularly with high parenteral dosage schedules, frequent evaluation of the renal and hematopoietic systems is recommended.

In suspected staphylococcus infections, proper laboratory studies (including sensitivity tests) should be performed.

The use of penicillin may be associated with the overgrowth of penicillin-insensitive organisms. In such cases, discontinue administration and take appropriate measures.

Adverse Reactions: Although serious allergic reactions are much less common with oral penicillin than with intramuscular forms, manifestations of penicillin allergy may occur.

Penicillin is a substance of low toxicity, but it possesses a significant index of sensitization. The following hypersensitivity reactions have been reported: skin rashes ranging from maculopapular eruptions to exfoliative dermatitis; urticaria; and reactions resembling serum sickness, including chills, fever, edema, arthralgia, and prostration. Severe and often fatal anaphylaxis has occurred (*see* Warnings). Hemolytic anemia, leukopenia, thrombocytopenia, and nephropathy are rarely observed side-effects and are usually associated with high parenteral dosage.

Administration and Dosage: Usual dosage range, 125 mg. (200,000 units) three times a day to 500 mg. (800,000 units) every four hours. For infants, 50 mg. per Kg. per day divided into three doses.

See package literature for detailed dosage instructions for prophylaxis of streptococcus infections, surgery, gonorrhea, and severe infections.

How Supplied: Tablets V-Cillin K[®] (Potassium Phenoxymethyl Penicillin Tablets, U.S.P.), 125 mg. (200,000 units), 250 mg. (400,000 units), and 500 mg. (800,000 units).

V-Cillin K[®] (potassium phenoxymethyl penicillin, Lilly), Pediatric, for Oral Solution, 125 mg. (200,000 units) and 250 mg. (400,000 units) per 5 cc. of solution (approximately one teaspoonful). [042567A] 800198

Additional information available to physicians upon request.

Eli Lilly and Company
Indianapolis, Indiana 46206



UNIQUE OFFICE RENTAL PLAN

at the

LEXINGTON PROFESSIONAL CENTER

133 EAST 73 STREET • NEW YORK, N. Y.

PAY ONLY FOR THE TIME & SERVICES USED !!

At the Lexington Professional Center you will have a handsomely furnished consultation room and fully equipped treatment room for **just the hours you need**—without any long term commitment, without any capital investment.

The modest cost will astonish you at this prestige medical center where you pay only for the hours and services you use. And you will have at your service the finest equipment and facilities . . . including X-Ray, radio-active isotope scanners, pharmacy *plus* 24 hour telephone answering, receptionist and secretarial services.

Full Time Offices Available, Furnished or Unfurnished

For additional information call Mrs. Ruth Freund, UN 1-9000

LEXINGTON PROFESSIONAL CENTER



133 E. 73rd ST., NEW YORK 21, N. Y.

UNiversity 1-9000

Over 4½ million prescriptions already filled at Macy's

*All 12 Macy Prescription Centres
satisfy your patients with
efficient professional service*



- You may phone in your patients' prescriptions (except for narcotics, amphetamines, and barbiturates).
- Your patients may charge their prescriptions on their Macy Shopping Accounts or C-T Plans
- Your patients may have their prescriptions delivered anywhere free of extra charge.
- We serve your patient with a full line of convalescent and diabetic needs.

Macy's Herald Square, OX 5-4400
Macy's Parkchester, TA 8-7000
Macy's Jamaica, OL 7-9000
Macy's Flatbush, UL 6-5000
Macy's Roosevelt Field, PI 6-8200
Macy's Huntington, AR 1-3000
Macy's White Plains, WH 6-5015
Macy's New Haven, Conn., 624-9271
Macy's Bay Shore, MO 5-8400
Macy's Queens, AR 1-9100
Macy's Colonie, 459-1950
Macy's New Rochelle, 633-7700

Five titles from
HAFNER PUBLISHING COMPANY

**HARTMAN, C. G. and
 STRAUS, W. L. (eds.)**

The Anatomy of the Rhesus Monkey

(Macaca Mulatta)

Illustrated by B. Kopel

ix, 383 pages, 128 illus. in the text and on
 plates. (1933) Reprint 1963 \$11.50
 B.C.L.

HERRICK, C. J.

**Neurological Foundations of
 Animal Behavior**

xii, 334 pages, 131 illus. (1924) Reprint
 with a new introduction, 1962 \$9.50

RAMON y CAJAL, S.

**Degeneration and Regeneration of the
 Nervous System**

Translated and edited by Dr. Raoul M. May
 2 vols., 769 pages, 317 illus. (1928)

Reprint 1968 \$25.00

THORNDIKE, EDWARD L.

Animal Intelligence

Experimental Studies

viii, illus., index, 297 pages. (1911)

Reprint 1965 \$5.25

McCLUNG, C. E. (ed.)

**Handbook of Microscopical Technique
 for Workers in Animal and Plant Tissues**

By 35 authors, 3rd revised edition by

R. McClung Jones.

xix, 790 pages, illus. (1950)

Reprint 1964

\$17.50

Order from

STECHERT-HAFNER, INC.

31 East 10th Street, New York, N. Y. 10003

Include MILK in your LOW-SALT DIETS

It's no longer necessary to deny patients fresh, fluid
 palatable Milk in low-salt diets. Walker-Gordon fresh
 Lo-Sodium Milk (Certified Milk with 90% of Sodium removed)
 contains less than 50 mg. Sodium per quart. *Guaranteed*
free of antibiotic residue. Paper half-pints for hospitals,
 quart bottles for home delivery. Write or phone for literature,
 low-sodium diet sheets, and professional sample.

Fresh
WALKER-GORDON LO-SODIUM MILK

Walker-Gordon Certified Milk Farm, Plainsboro, N. J. ★ (609) 799-1234

New York: (212) Walker 5-7464 ★ Phila.: (215) MA 7-6338


Also Certified Raw, Homogenized Vitamin D and Skimmed Milks;
 available through leading Milk Dealers or call Walker-Gordon

GREAT AMERICANS WHO WERE DOCTORS...

NUMBER ONE OF A SERIES



Josiah Bartlett
(1729-1795)



Matthew Thornton
(c1714-1803)

Aside from the active practice of medicine, these two physicians were also signatories of the Declaration of Independence. Dr. Bartlett enhanced his reputation in the mid-1750's by his novel treatment (with Peruvian bark) of the angina maligna. His career was distinguished by his term as Chief Justice of the Supreme Court in 1788.

WE AT GHI came along too late — by about 200 years — to be involved with the Declaration of Independence. We do, however, see our role in the community as an important one. Why? Because we are innovators. Over the years we have added many types of protection to our covered benefits — such as home calls, office visits, and ambulatory psychiatric care. By bringing doctor care to people who might otherwise have gone without until it was too late, we have strengthened the fabric of the society of which we are a part. By becoming a GHI Participating Family Doctor or a GHI Participating Physician, you can become part of the team that performs these worthwhile services.

Join the approximately 10,000 doctors who now participate in GHI's Plans. Write to us today for more information about why and how you should participate in GHI. Do it now — while you're thinking about it!



Group Health Insurance, Inc. / 227 West 40th Street, New York, N.Y. 10018 / Phone: 564-8900

Whenever anxiety induces or intensifies clinical symptoms

Librium®

(chlordiazepoxide HCl)

*Quickly relieves anxiety ~ Helps improve response
in psychophysiologic disorders ~ Seldom
impairs mental acuity or physical coordination,
on proper dosage ~ Has wide margin of safety*

Before prescribing, please consult complete product information, a summary of which follows:

Indications: Indicated when anxiety, tension and apprehension are significant components of the clinical profile.

Contraindications: Patients with known hypersensitivity to the drug.

Warnings: Caution patients about possible combined effects with alcohol and other CNS depressants. As with all CNS-acting drugs, caution patients against hazardous occupations requiring complete mental alertness (e.g., operating machinery, driving). Though physical and psychological dependence have rarely been reported on recommended doses, use caution in administering to addiction-prone individuals or those who might increase dosage; withdrawal symptoms (including convulsions), following discontinuation of the drug and similar to those seen with barbiturates, have been reported. Use of any drug in pregnancy, lactation, or in women of childbearing age requires that its potential benefits be weighed against its possible hazards.

Precautions: In the elderly and debilitated, and in children over six, limit to smallest effective dosage (initially 10 mg or less per day) to pre-

clude ataxia or oversedation, increasing gradually as needed and tolerated. Not recommended in children under six. Though generally not recommended, if combination therapy with other psychotropics seems indicated, carefully consider individual pharmacologic effects, particularly in use of potentiating drugs such as MAO inhibitors and phenothiazines. Observe usual precautions in presence of impaired renal or hepatic function. Paradoxical reactions (e.g., excitement, stimulation and acute rage) have been reported in psychiatric patients and hyperactive aggressive children. Employ usual precautions in treatment of anxiety states with evidence of impending depression; suicidal tendencies may be present and protective measures necessary. Variable effects on blood coagulation have been reported very rarely in patients receiving the drug and oral anticoagulants; causal relationship has not been established clinically.

Adverse Reactions: Drowsiness, ataxia and confusion may occur, espe-

cially in the elderly and debilitated. These are reversible in most instances by proper dosage adjustment, but are also occasionally observed at the lower dosage ranges. In a few instances syncope has been reported. Also encountered are isolated instances of skin eruptions, edema, minor menstrual irregularities, nausea and constipation, extrapyramidal symptoms, increased and decreased libido—all infrequent and generally controlled with dosage reduction; changes in EEG patterns (low-voltage fast activity) may appear during and after treatment; blood dyscrasias (including agranulocytosis), jaundice and hepatic dysfunction have been reported occasionally, making periodic blood counts and liver function tests advisable during protracted therapy.

Usual Daily Dosage: Individualize for maximum beneficial effects. *Oral*—Adults: Mild and moderate anxiety and tension, 5 or 10 mg t.i.d. or q.i.d.; severe states, 20 or 25 mg t.i.d. or q.i.d. Geriatric patients: 5 mg b.i.d. to q.i.d. (See Precautions.)

Supplied: Librium® (chlordiazepoxide HCl) Capsules, 5 mg, 10 mg and 25 mg—bottles of 50. LibritabsTM (chlordiazepoxide) Tablets, 5 mg, 10 mg and 25 mg—bottles of 100. With respect to clinical activity, capsules and tablets are indistinguishable.



Roche®

LABORATORIES

Division of Hoffmann-La Roche Inc.
Nutley, New Jersey 07110

Also available:
LibritabsTM (chlordiazepoxide)
5-mg, 10-mg, 25-mg tablets